

Note: Various size labels
attached below

Doc. ID 588358

ACCEPTED
FOR REGISTRATION
Aug. 01, 2023

New York State Department
of Environmental Conservation
Division of Solid & Hazardous Materials
Pesticide Product Registration

SA SOUTHERN
50 AG®

**MAKES UP TO
24 GALLONS**

LIQUID COPPER FUNGICIDE

ACTIVE INGREDIENT _____ **BY WT.**
Copper diammonia diacetate*..... 27.15%
complex.*

[Bis(acetate-O) Diamminecopper]

CAS No. 13822-80-5

OTHER INGREDIENTS:..... 72.85%

Total..... 100.00%

*Metallic Copper Equivalent, 8.0%

Contains 0.04825 LBS Copper per 8 ounces

FOR RESIDENTIAL
USE ONLY

**Controls Many
Diseases Attacking
Citrus, Vegetables,
Ornamentals,
Fruits & Algae in
Ornamental Lawns
and Turf.**



**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See label booklet for continued
Precautionary Statements and
Directions for use.


Net Contents Liquid: 8 ounces (.237 liters)

Distributed by: **Southern**

Agricultural Insecticides, Inc.
Palmetto, FL 34220 Hendersonville, NC 28793

Boone, NC 28607 FZ

EPA Reg. No. 10465-3-829 EPA Est. No. 829-FL-1



0 51538 02901 8

S A SOUTHERN
5 0 AG®

LIQUID COPPER FUNGICIDE

FOR RESIDENTIAL USE ONLY

**Controls Many Diseases on Ornamentals, Vegetables,
Fruits, Citrus, & Algae in Turfgrass.**

<u>ACTIVE INGREDIENT</u>	<u>BY WT.</u>
Copper diammonia diacetate* complex.*	27.15%
[Bis(acetate-O) Diamminecopper]	
CAS No. 13822-80-5	
INERT INGREDIENTS:.....	72.85%
Total.....	100.0%

*Metallic Copper Equivalent, 8.0%

Contains 0.1935 LBS Copper per quart




CAUTION
KEEP OUT OF REACH
OF CHILDREN

See inside of booklet for continued Precautionary
Statements and Directions for use.

Net Contents:
1 Quart (.946 liters)

Distributed by: **Southern**
Agricultural Insecticides, Inc.
Palmetto, FL 34420 Hendersonville, NC 28793
Boone, NC 28607 F3
EPA Reg. No. 10465-3-829 EPA Est. No. 829-FL-1



SA SOUTHERN
50 AG®

**MAKES UP
TO
48 GALLONS**

LIQUID COPPER FUNGICIDE

FOR RESIDENTIAL USE ONLY

**Controls Many Diseases on Ornamentals, Vegetables,
Fruits, Citrus, & Algae in Turfgrass.**

<u>ACTIVE INGREDIENT</u>	<u>BY WT.</u>
Copper diammonia diacetate*.....	27.15%
complex.*	
[Bis(acetate-O) Diamminecopper]	
CAS No. 13822-80-5	
INERT INGREDIENTS:.....	72.85%
Total.....	100.0%
*Metallic Copper Equivalent, 8.0%	
Contains 0.0965 LBS Copper per pint	




KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside of booklet for continued Precautionary
Statements and Directions for use.

**Net Contents
1 Pint (.474 liters)**

Distributed by: **Southern
Agricultural Insecticides, Inc.**
P.O. Box 218 Palmetto, FL 34220
F2 941-722-3285
EPA Reg. No. 10465-3-829 EPA Est. No. 829-FL-1



FRONT OF BOOKLET

LIQUID COPPER FUNGICIDE

FOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT	BY WT.
Copper diammonia diacetate complex*	27.15%
[Bis(acetate-O) Diamminecopper]	
CAS No. 13822-80-5	
<u>OTHER INGREDIENTS:.....</u>	<u>72.85%</u>
Total.....	100.00%

*Metallic Copper Equivalent, 8.0%
Contains 0.772 lbs Copper per gallon

*See inside of booklet for First Aid,
continued Precautionary Statements, and
Directions for use.*

CAUTION
KEEP OUT OF REACH OF CHILDREN

Distributed by:
Southern Agricultural Insecticides, Inc.
P.O. Box 218 Palmetto, FL 34220
EPA REG. NO. 10465-3-829
EPA EST. NO. 829-FL-1 B4A

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OF CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER: Have product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes eye and skin irritation. Harmful if swallowed, absorbed through the skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wear protective eyewear (goggles, face shield or safety glasses), long sleeved shirt and long pants, shoes and socks and chemical-resistant gloves when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse. Do not use food utensils such as teaspoons or tablespoons for food purposes after use with pesticides.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Only protected users may be in the area during application. Do not enter or allow others to enter the treated area until sprays have dried.

Shake well before using.

INFORMATION

The control of diseases with fungicides is based on PREVENTION plant surfaces must be completely covered with the fungicide to successfully prevent infection. Copper pesticides are considered preventative, not curative of plant diseases. Use the highest indicated rate per crop when disease incidence is high or expected to be, depending on rainfall and temperature. The lower rate is suitable for general preventive sprays under normal conditions. Since weather conditions and disease incidence vary, consult your Agricultural Extension Service for timing and initial application.

PRODUCT INSTRUCTIONS

Seek advice on identifying specific disease problems and solutions from your local Agricultural Extension Agent or other knowledgeable persons.

Partially fill the spray tank/container with water, add the desired amount of Liquid Copper Fungicide and continue filling the tank/container. If applied with other products, add Liquid Copper Fungicide last. Agitate the tank during mixing and application, until tank is empty. A plastic or metal stirring stick should provide adequate agitation. Observe all cautions and limitations on labeling of all products used in mixtures. Start with clean equipment. Equipment should be flushed well with water after use.

APPLICATION

Fruit: Mix specified rate in 1 gallon of water and apply 3 gallons of mixed solution to a small tree or bush, 6 gallons of mixed solution to a medium size tree, or 9 gallons of mixed solution to a large tree. Thoroughly spray tree to point of runoff, including upper and lower surfaces of foliage. Do not overspray. Do not mix more spray solution than needed.

Vegetables: Mix specified rate in 1 gallon of water and apply 2 gallons of mixed solution per 1,000 sq. ft. (1 gallon of mixed solution per 500 sq. ft.)

Miscellaneous: For trees: Mix specified rate in 1 gallon of water and apply 3 gallons of mixed solution to a small tree, 6 gallons of mixed solution to a medium size tree, or 9 gallons of mixed solution to a large tree. Thoroughly spray tree to point of runoff, including upper and lower surfaces of foliage. Do not overspray. Do not mix more spray solution than needed.

NOTE: This product may be reactive on metal and masonry surfaces such as galvanized roofing. AVOID contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

FRUITS

Fruit	Diseases Controlled	Rate/Gallon of Water	Instructions
Apples	Anthracnose	4-5 tsp.	Apply as a dormant spray once to foliage after harvest.
	Apple scab (black spot) Bacterial canker Blossom and shoot blast	4-6 tsp.	Apply as a dormant spray once post-harvest before fall rains.
	Fireblight	1/2- 1 tsp.	Apply at 10% bloom and repeat at no less than 5-7 day intervals during the bloom period. Do not use on copper-sensitive varieties.
			4-6 tsp.
NOTE: Do not apply more than 370 tsp. of Liquid Copper Fungicide (0.37 lbs. metallic copper) to Apples per 1,000 sq. ft. per year.			
Avocados	Anthracnose	4 tsp.	Apply when the flower buds begin to swell and continue at monthly intervals until August.
NOTE: Do not apply more than 430 tsp. of Liquid Copper Fungicide (0.43 lbs. metallic copper) to Avocados per 1,000 sq. ft. per year.			

Fruit	Diseases Controlled	Rate/Gallon of Water	Instructions
Bananas	Sigatoka	1-2 tsp.	Apply every 3-4 weeks.
	Black pitting	2.5 fl. oz.	Mix in 1 gallon of water. Apply directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence. Do not apply at less than 7 day intervals

NOTE: Do not apply more than 71.6 fl. oz. of Liquid Copper Fungicide (430 tsp./0.43 lbs. metallic copper) to Bananas per 1,000 sq. ft. per year.

Blueberries	Bacterial Canker	2-5 tsp.	Apply before fall rains and again 4 weeks later.
	Cane Canker	2-5 tsp.	Apply before fall rains and again 4 weeks later. In the spring during wet weather, apply at 10-14 day intervals beginning at leaf emergence.

NOTE: Do not apply more than 190 tsp. of Liquid Copper Fungicide (0.19 lbs. metallic copper) to Blueberries per 1,000 sq. ft. per year.

Citrus	Greasy Spot Melanose Pink Pitting Scab	2-4 tsp.	Apply as pre-bloom and post bloom sprays. Do not apply at less than 7 day intervals
	Brown rot	1-3 tsp.	Apply in the fall before or just after heavy rains. In areas of skirt sprays, apply to a height of at least 4 feet.

(Citrus continued on next column)

Fruit	Diseases Controlled	Rate/Gallon of Water	Instructions
(Citrus continued)			(Skirt sprays are applications up to 4 ft. high on the tree. The whole tree is not sprayed.) Do not apply at less than 7 day intervals.
NOTE: Do not apply more than 290 tsp. of Liquid Copper Fungicide (0.29 lbs. metallic copper) to Citrus per 1,000 sq. ft. per year.			
Limes	Greasy spot	4 tsp.	Apply in June and continue at monthly intervals through August.
NOTE: Do not apply more than 290 tsp. of Liquid Copper Fungicide (0.29 lbs. metallic copper) to Limes per 1,000 sq. ft. per year.			
Mangos (NOT FOR USE IN CALIFORNIA)	Anthracnose	4 tsp.	Apply at 30 day intervals from the time the panicles are 2 inches high until all fruits are set until August
NOTE: Do not apply more than 145 tsp. of Liquid Copper Fungicide (0.145 lbs. metallic copper) to Mangos per 1,000 sq. ft. per year.			
Peaches	Bacterial spot	4 tsp.	Apply as a dormant spray at no less than 7 day intervals.
Nectarines	Blossom Brown Rot Leaf Curl Shot Hole	4-6 tsp.	Apply as a dormant and delayed dormant spray to protect buds and shoots from infection during rainy period. Reapply up until late bud swell. Do not apply at less than 7 day intervals. Do not apply after full bloom.
NOTE: Do not apply more than 410 tsp. of Liquid Copper Fungicide (0.41 lbs. metallic copper) to Peaches/Nectarines per 1,000 sq. ft. per year.			

Fruit	Diseases Controlled	Rate/Gallon of Water	Instructions
Strawberries	Leaf Spot Scorch	3-4 tsp.	Apply at 7-10 day intervals from the time new growth starts until harvest.

NOTE: Do not apply more than 190 tsp. of Liquid Copper Fungicide (0.19 lbs. metallic copper) to Strawberries per 1,000 sq. ft. per year.

*Delayed Dormant - The period just before blossom buds break open.

VEGETABLES

Vegetables	Diseases Controlled	Rate/Gallon of Water	Instructions
Beans, Peas, Lentils (Succulent and Dry)	Bacterial Blight (Halo and Common)	2-6 tsp.	Apply when plants are 3-5 inches high and before diseases appear. Repeat at 7-10 day intervals. Apply at 7 day intervals if disease pressure is severe.

NOTE: Do not apply more than 30 tsp. of Liquid Copper Fungicide (0.03 lbs. metallic copper) to Beans/Peas/Lentils per 1,000 sq. ft. per year.

Crucifers [Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Greens (Collard, Mustard and Turnip)]	Blackleaf Spot Black Rot	2-6 tsp.	Apply when disease appears and repeat at 7-10 day intervals.
	Downey Mildew	1-2 tsp.	Apply when disease appears and repeat at 7-10 day intervals.
	NOTE: Do not apply more than 60 tsp. of Liquid Copper Fungicide (0.06 lbs. metallic copper) to Crucifers per 1,000 sq. ft. per year.		

Vegetables	Diseases Controlled	Rate/Gallon of Water	Instructions
Cucurbits (Cantaloupe, Cucumber, Honeydew Muskmelon Pumpkin, Squash Watermelon)	Alternaria Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Gummy Stem Blight Powdery Mildew Watermelon Bacterial Fruit Blotch	3-4 tsp.	Apply when disease appears and repeat 7-10 day intervals.
NOTE: Do not apply more than 115.2 tsp. of Liquid Copper Fungicide (0.12 lbs. metallic copper) to Cucurbits per 1,000 sq. ft. per year.			
Eggplant	Alternaria Blight Anthracnose Phomopsis	4 tsp.	Apply before disease appears and repeat at 7-10 day intervals.
NOTE: Do not apply more than 180 tsp. of Liquid Copper Fungicide (0.18 lbs. metallic copper) to Eggplant per 1,000 sq. ft. per year.			
Onions	Downy Mildew Purple Blotch	4 tsp.	Apply when plants are 4-6 inches high and repeat at 7-10 day intervals.
NOTE: Do not apply more than 140 tsp. of Liquid Copper Fungicide (0.14 lbs. metallic copper) to Onions per 1,000 sq. ft. per year.			
Peppers	Bacterial Spot Cercospora Leaf Spot	3-6 tsp.	Make first application upon emergence of seedlings or immediately after transplanting and repeat at 7-10 day intervals.

(Peppers continued on next column)

Vegetables	Diseases Controlled	Rate/Gallon of Water	Instructions
Peppers (continued)			When disease is severe, apply at 7 day intervals. NOTE: Disease control is critical during fruiting.
NOTE: Do not apply more than 270 tsp. of Liquid Copper Fungicide (0.27 lbs. metallic copper) to Peppers per 1,000 sq. ft. per year.			
Potatoes	Early Blight Late Blight	3-6 tsp.	Apply on first appearance of disease and repeat at 7-10 day intervals.
NOTE: Do not apply more than 570 tsp. of Liquid Copper Fungicide (0.57 lbs. metallic copper) to Potatoes per 1,000 sq. ft. per year.			
Spinach	Anthraxnose Cercospora Leaf Spot Downy Mildew	3 tsp.	Apply on first appearance of disease and repeat at 7-10 day intervals.
NOTE: Do not apply more than 90 tsp. of Liquid Copper Fungicide (0.09 lbs. metallic copper) to Spinach per 1,000 sq. ft. per year.			
Tomatoes	Bacterial Speck Bacterial Spot Early Blight Late Blight	3-5 tsp.	Make first application upon emergence of seedlings or immediately after transplanting and repeat at 7-10 day intervals. When disease pressure is severe, apply at the shorter spray interval. Complete coverage is essential for disease control.

(Tomatoes instructions continued on next page)

(Tomatoes instructions continued) NOTE: While the labeled rate is particularly effective against Bacterial Spot, a tank mix of Maneb or Mancozeb used at the labeled rates controls a broad range of diseases.

NOTE: Do not apply more than 400 tsp. of Liquid Copper Fungicide (0.4 lbs. metallic copper) to Tomatoes per 1,000 sq. ft. per year.

MISCELLANEOUS

To control Ball Moss in Live Oak, apply 2 fl. oz. per gallon of water. Apply in the spring when ball moss is actively growing using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. A second application may be required after 12 months. NOTE: Liquid Copper Fungicide may be injurious to ornamentals grown under Live Oaks. This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

NOTE: Do not apply more than 76.7 fl. oz. of Liquid Copper Fungicide (0.46 lbs. metallic copper) to Live Oak per 1,000 Sq. ft. per year.

TURFGRASS

To control algae in ornamental lawns and turf, apply 10 fl. oz. (0.06 lbs. metallic copper) of Liquid Copper Fungicide in 5 gallons of water per 1,000 square feet. Liquid Copper Fungicide may be used alone or in combination with other registered fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes. Do not apply more than 80.0 fl. oz. of Liquid Copper Fungicide (0.48 lbs. metallic copper) per 1,000 sq. ft. per year. Do not apply at less than 10 day intervals.

NOTE: Phytotoxicity may occur depending upon varietal differences. Apply the directed rate to a small area and observe for 7 to 10 days for signs of injury. If phytotoxicity occurs, discontinue use. Do not apply in a spray solution with a pH of less than 6.5.

ORNAMENTALS

For use in container, landscape and outdoor ornamental plantings. Liquid Copper Fungicide can also be used on ornamentals in home greenhouses and shadehouses.

Notice to User: Plant sensitivities to Liquid Copper Fungicide have been found to be acceptable in specific genera and species listed on this label; however, it is impossible to know sensitivities under all conditions and phytotoxicity may occur. Due to the large numbers of species and varieties of ornamentals, it is impossible to test every one for sensitivity to Liquid Copper Fungicide. Neither the manufacturer nor seller recommends use upon species not listed on the label nor has it been determined that Liquid Copper Fungicide can safely be used on ornamental plants not listed on this label. The user should determine if Liquid Copper Fungicide can be used safely prior to use.

Apply as a thorough coverage spray using 2 teaspoons of Liquid Copper Fungicide per gallon of water. Begin application at first sign of disease and repeat at 7-14 day intervals as needed. Use shorter interval during periods of frequent rains or when severe disease conditions persist. Do not apply more than 460 tsps. of Liquid Copper Fungicide (0.46 lbs. metallic copper) per 1,000 sq. ft. per year.

Liquid Copper Fungicide may be used alone or in combination with other registered fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes. NOTE: Do not tank mix Liquid Copper Fungicide with Aliette fungicide unless appropriate precautions have been taken to buffer the spray solution. Severe phytotoxicity may result if adequate precautions are not taken.

PLANT	DISEASE
Althea (Rose of Sharon)	Bacterial leaf spot
Aralia	Alternaria, Cercospora leaf spot
Arborvitae	Alternaria twig blight, Cercospora leaf blight
Azalea (1)	Botrytis blight, Cercospora leaf spot, Phytophthora dieback, Powdery mildew
Begonia	Bacterial leaf spot (Erwina sp., Pseudomonas sp., Xanthomonas sp.)
Bougainvillea	Anthracnose, Bacterial leaf spot
Bulbs (Tulip, Gladiolus)	Anthracnose, Botrytis blight
Camellia	Anthracnose, Bacterial leaf spot
Camphor tree	Pseudomonas leaf spot

PLANT**DISEASE**

Canna	Pseudomonas leaf spot
Carnation (1)	Alternaria blight, Pseudomonas leaf spot, Botrytis blight
Chinese tallow tree	Bacterial leaf spot (Xanthomonas sp., Pseudomonas sp.)
Chrysanthemum (1)	Septoria leaf spot, Botrytis blight
Cotoneaster	Botrytis blight
Dahlia	Alternaria leaf spot, Botrytis gray mold, Cercospora leaf spot
Date Palm	Pestalotia leaf spot
Dianthus	Bacterial spot, Bacterial soft rot
Dogwood	Anthracnose
Dusty Miller	Bacterial leaf spot (Pseudomonas cichorii)
Echinacea	Bacterial leaf spot (Pseudomonas cichorii)
Elm "Drake"	Xanthomonas leaf spot
Euonymus	Botrytis blight, Anthracnose
European fan palm	Pestalotia leaf spot
Gardenia	Alternaria leaf spot, Botrytis bud rot, Cercospora leaf spot
Geranium	Alternaria leaf spot, Botrytis gray mold, Cercospora leaf spot
Gladiolus	Alternaria leaf spot, Botrytis gray mold, Bacterial leaf blight
Goldenrain tree	Bacterial leaf spot
Hibiscus	Bacterial leaf spot
Holly fern	Pseudomonas leaf spot
Impatiens	Bacterial leaf spot
India hawthorn (2)	Anthracnose, Entomosporium leaf spot
Ivy (English, Algerian) (1)	Xanthomonas leaf spot
Ixora	Xanthomonas leaf spot
Juniper (Eastern red cedar)	Anthracnose
Lantana	Bacterial leaf spot
Lilac	Cercospora leaf spot
Loblolly bay	Anthracnose

PLANT	DISEASE
Loquat	Entomosporium maculata, Colletotrichum sp.
Magnolia (Saucer)	Bacterial leaf spot
Magnolia (Southern)	Algal leaf spot, Anthracnose, Bacterial leaf spot
Magnolia (Sweet bay)	Anthracnose
Mandevillas	Anthracnose
Marigold	Alternaria leaf spot, Botrytis leaf and flower rot, Seersucker leaf spot
Mulberry, weeping	Bacterial leaf spot
Oak, laurel	Algal leaf spot (Cephaleuros virescens)
Oleander	Bacterial leaf spot, Fungal leaf spot
Pachysandra	Volutella leaf blight
Pansy	Downy mildew
Pear (Flowering)	Fireblight, leaf spot
Pentas (Egyptian star)	Bacterial leaf spot (Xanthomonas sp.)
Peony	Botrytis blight
Periwinkle	Phomopsis stem blight
Philodendron	Bacterial leaf spot
Phlox	Alternaria leaf spot
Photinia	Anthracnose, Entomosporium
Pistachio	Anthracnose
Plantain lily	Bacterial leaf spot
Powder puff plant	Bacterial leaf spot
Pyracantha	Fireblight, scab
Queen palm	Exosporium leaf spot, Phytophthora bud rot
Rhododendron	Alternaria flower spot
Rose (1)	Powdery mildew, Black spot
Verbena	Xanthomonas leaf spot
Viburnum	Anthracnose
Washingtonia palm	Pestalotia leaf spot

PLANT**DISEASE**

Weeping willow

Anthracnose

Yucca (Adam's needle)

Cercospora leaf spot, Septoria leaf spot

(1) Discoloration of foliage and/or blooms has been noted on some varieties.

(2) For India Hawthorn use 2-4 level tablespoons per gallon.

FROST INJURY PROTECTION

Bacterial Ice Nucleation Inhibitor - Application of Liquid Copper Fungicide made to all plants listed on this label at rates indicated, just prior to anticipated frost conditions, will sustain control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia Herbicola* and *Pseudomonas fluorescence*) and may therefore provide protection against light frost.

Not recommended for those geographic areas where weather conditions favor severe frost.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a secure, locked area away from pets and out of the reach of children. Store only in original container and place in a locked storage area.

PESTICIDE DISPOSAL/CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE

Read the entire Directions for use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the

unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of the product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use of application, all of which are beyond the control of Southern Agricultural Insecticides, Inc. To the extent permitted by applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Southern Agricultural Insecticides, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Southern Agricultural Insecticides, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Southern Agricultural Insecticides, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Southern Agricultural Insecticides, Inc.'s election, the replacement of product.